

From Leningrad to the Day-Care Center The Ubiquitous *Giardia lamblia*

WAYNE X. SHANDERA, MD, Houston

Giardiasis is recognized as a worldwide public health problem. Seroprevalence data from both the developing and developed world show high rates of carriage in populations at risk for fecal-oral transmission, such as children in day-care centers. Outbreak investigation has expanded our understanding of reservoirs for *Giardia lamblia* and of the routes of transmission. Various host factors have been associated with infection. The pathogenesis of giardial infections is being elucidated, in particular the role of lectin activation in producing disease. Three standard chemotherapeutic agents are available in the United States. The institution of community-wide prevention measures is equally important. Current areas of investigation including antigenic composition and enzymatic variants should result in effective forms of immunotherapy, while more effective forms of chemoprophylaxis could assist in eradicating the pathogen from institutional settings.

(Shandera WX: From Leningrad to the day-care center—The ubiquitous *Giardia lamblia*. West J Med 1990 Aug; 153:154-159)

A description of the pathogen responsible for giardiasis may be found in the works of Antony van Leeuwenhoek, who discovered in 1681:

animalcules . . . all of the one and the same make; their bodies [are] somewhat longer than broad and their bellies [are] flatlike, furnished with sundry little paws, wherewith they made such a stir in the clear medium, and among the globules, that you might e'en fancy you saw a pissabed [woodlouse] running up against a wall; and albeit they made a quick motion with their paw, yet for all that they made but small progress.¹

Such "animalcules," probably *Giardia lamblia*, were not subsequently described until a Bohemian physician, Wilhelm Lambl, ascribed to the organism its first scientific name, *Cercomonas intestinalis*, and elucidated its place in the ecosystem, including it among the fecal flora.² Western Europeans recognize his contribution in one of their two names for the pathogen: *Lamblia intestinalis*³ or *Giardia intestinalis*. The term *Giardia* appears in discussions written by investigators from Berkeley, California, and Germany during the early 20th century; *Giardia* honors a French biologist named Alfred Giard. Hence, the name *Giardia lamblia* honors two 19th-century European scientists.^{4,5}

With the modern growth in travel and increased attention to community epidemiology, the importance of *Giardia* species in human disease resurfaced. In the 1960s Americans returning from the Soviet Union with watery diarrhea carried the pathogen in stool; investigations pinpointed the water system in Leningrad.⁶ A reported community-wide outbreak wherein *G lamblia* was incriminated occurred in Aspen in 1964-1965: protracted episodes of gastroenteritis developed among those who drank from the city water supply.⁷ Such outbreak investigations have supplemented studies of antigenic composition, ultrastructure, life cycle, and enzyme variants in contributing to understanding *G lamblia* and giardiasis.

The Organism and Its Epidemiology

The life cycle of this flagellated protozoan is relatively simple: after cysts are ingested by a mammal, gastric acidity induces excystment and trophozoite development.⁸ Trophozoites cause disease, although the mechanism by which disease occurs is debated. Organisms reencyst in the intestine on bile salt exposure⁹ and excretion of cysts ensues,

although cyst excretion does not have the strict disease dependence of trophozoite excretion, trophozoites being found only in diarrheic stools. The incubation period is usually between 7 and 15 days but may be much longer.¹⁰ The trophozoites are very labile outside the host; cysts survive outside the host if not dehydrated or subjected to extreme temperatures.¹¹ The estimated infective dose in 50% of the population is between 10 and 100 cysts.¹²

Giardia species infections are ubiquitous. Outbreaks in the developed world have been reported from North America, Europe, and Australia¹³⁻²⁰; in the developing world, high rates of endemicity have been found in Bangladesh,²¹ Nigeria,²² Guatemala,²³ and Peru.²⁴ The World Health Organization estimates that about 200 million infections occur annually in the developing world.²⁵

In most communities the rate of endemic disease is unknown: giardiasis is often not reportable, variations exist in the methods of collecting specimens, and medical personnel vary in diagnostic skills. In the United States, 4% of stool specimens submitted to state health department laboratories are positive for *Giardia* species.²⁶ In one jurisdiction with required reporting, the incidence was 17.5 per 100,000 per year; 50% of cases occurred in day-care centers (SW Washington Health District, unpublished data, September 1985).

Community-wide outbreaks are typically from a common source. Although food and body fluids are recognized vehicles of transmission, water is the most common vehicle: the Public Health Service reports that 12 of 99 waterborne outbreaks during a four-year interval were due to *G lamblia*.^{27,28} Camping-associated outbreaks are caused by ingesting contaminated water.^{11,29} A community swimming pool-associated outbreak may have been due to a toddler soiling in the pool,³⁰ and a swimming slide-associated outbreak may have developed similarly.³¹ Food outbreaks are uncommon; a Minnesota outbreak was associated with ingesting home-canned salmon prepared by an infected handler and served at a rural school³²; a second in Connecticut was associated with cold noodle salad.³³

Important community outbreaks include one in Camas, Washington (suburban Portland, Oregon), during 1976 that

ABBREVIATIONS USED IN TEXT

AIDS = acquired immunodeficiency syndrome
IFA = indirect fluorescent antibody

affected 128 persons when the surface water supply broke down. Chlorinated and filtered water supplies failed to prevent the outbreak, and water treatment required flocculation, sedimentation, and coagulation.¹³ A 1980 outbreak in Red Lodge, Montana, showed how climatic and environmental factors interact in producing disease. After the Mount Saint Helens (Washington) eruption, the deposit of volcanic ash that had been blown eastward, in conjunction with increased snowmelt associated with unusually warm weather, raised the city water supply level; the elevation provided animals access to surface water.¹⁴ Ski resorts have been susceptible to *Giardia* species outbreaks, with reports from both Vail and Aspen, Colorado^{7,15}; attack rates in Aspen were associated with the amount of water ingested and the length of residence—longer-term residents in 1981 had lower attack rates, ostensibly because of exposure during a 1964 outbreak.¹⁵ A 1978 New England outbreak revealed the asymptomatic nature of many community-wide infections: 75% of those infected did not require treatment.¹⁶

Intercontinental travel and migration have been studied in East Germany: foreigners arriving from tropical areas showed higher rates of seroprevalence to *G lamblia* than did German travelers to and from the tropical areas, who had higher rates than German locals.³⁴ Other studies have outlined the high rates of giardiasis among immigrants from Third World countries.³⁵

Well-substantiated methods of transmission also include sexual transmission and person-to-person. Three sexually transmitted disease centers have reported *G lamblia* among the fecal flora of homosexual and bisexual men.³⁶⁻³⁸ In a case-control study, 18% of 126 homosexual men carried *Giardia* species, compared with 2% of inpatient and outpatient controls at New York Hospital.³⁹ Oral-anal sex, however, is not as established a predisposing factor toward giardiasis as it is toward amebiasis and helminthic infections.

Day-care center transmission has been substantiated in the United States, Canada, and Australia⁴⁰⁻⁴⁵; age-specific

attack rates confirm that childhood disease maintains *Giardia* species in the community. In data from Houston, between 21% and 26% of 600 attendants at 30 day-care centers excreted *Giardia* cysts (trophozoite excretion between 3% and 4%).⁴⁰ Case-control methods have revealed that age-matched children not in day-care centers have lower isolation rates than those in day care: 2% versus 29% to 54%.⁴¹ Australian investigators emphasize high rates of asymptomatic infection among children.⁴² Family members and household contacts of infected children are at risk, secondary infection rates being between 17% and 47%.^{40,42-45}

In summary, the epidemiology of *G lamblia* reveals that transmission occurs by many methods—common sources, sexual transmission, person-to-person nonsexual contact—but water is the usual source of community-wide outbreaks. Environmental factors that may be determinants of outbreaks include the presence of infected animals, the availability of surface water, and climate factors influencing the water supplies. A high prevalence of infection among toddlers and infants is a consequence of fecal-oral transmission in the developed world and water and food exposures in the developing world.

Host Risk Factors

The prevalence of giardiasis, as with any infection, is dependent on host, agent, and environmental factors. Environmental factors have been mentioned; agent factors will be discussed in another section. Host risk factors include sex- and age-specific attack rates. Males are infected more commonly than females.⁴⁶⁻⁵¹ Age-specific prevalence data show a propensity toward infection in childhood; the prevalence of infection in the younger-than-5 cohort ranges between 4% and 42% (Table 1).^{21-23, 46-54} The highest rates occurred in the Nile delta where among 2- to 4-year-old children, 42% showed cyst or trophozoite positivity, with excretion lasting as long as 17 weeks.⁵⁴ A difficulty establishing an association between infection and symptomatic illness is a common theme; studies from Peru report frequent reinfections after treatment.²⁴ Other possible host risk factors include immunodeficiency, human leukocyte antigen phenotype, nutrition status, achlorhydria, blood type, and concomitant infection with other agents, such as *Cryptosporidium* species or *Candida albicans*.

TABLE 1.—Childhood Prevalence of *Giardia* Species Stool Carriage

Population of Endemicity, Age	Children, No.	Stool Specimens Examined, No.	Methods	Symptomatic, %	Prevalence, %		Reference Source
					Childhood	All Ages	
Hawaiian school children.....	390	1	Wet mount	0	4-5	..	Desowitz and Wiebenga, 1975 ⁴⁶
Nîmes, France.....	> 6	..	Fontaine et al, 1984 ⁴⁷
Ontario day-care attendees, 2-5 years	244	1	Zinc sulfate	0	6-8	..	Woo and Paterson, 1986 ⁴⁸
Washington state infants, 1-3 years	518	1	FEA	..	7.1*	..	Harter et al, 1982 ⁴⁹
Philadelphia inner-city children.....	385	..	PVA-FE	..	8-20	..	Weiner et al, 1959 ⁵⁰
Northwestern Australian children, 1-5 years..	50	1	PVA-zinc sulfate	32†	12	4	Boreham and Phillips, 1986 ⁵¹
Nigerian children in endemic area, 1-5 years .	449	1-3	FSE/iodine	..	13.8	7	Oyerinde et al, 1977 ²²
Rural Somali children and mothers	517	1	Formalin/FE	..	15-16	..	Peltola et al, 1988 ⁵²
Guatemalan children, 0-3 years.....	45	Weekly	PVA/FE/trichrome	0-8.2	20.2	..	Farthing et al, 1986 ²³
Costa Rican children younger than 5 years...	293	1	..	‡	25.3	..	Moore et al, 1966 ⁵³
Bangladeshi children younger than 5 years...	91	..	Iodine	..	21-51§	..	Gilman et al, 1985 ²¹
Rural Nile delta children, 2-4 years.....	42	Weekly	Trichrome	..	42.4	17	Sullivan et al, 1988 ⁵⁴

FE = formalin-ether, FEA = formalin-ethyl acetate, FSE = formalin-saline-ether, PVA = polyvinyl alcohol

*21.3% among family member contacts.

†By history.

‡No difference in isolation rate was seen in the presence of diarrhea.

§Seroprevalence is higher among malnourished patients.

||Cysts and trophozoites.

A variety of immunodeficient states—common, variable; Bruton's; the acquired immunodeficiency syndrome (AIDS)—have been studied with respect to giardiasis.^{55,56} Persons with immunodeficiencies may not be at particular risk for severe giardiasis developing.⁵⁷⁻⁵⁹ X-linked (Bruton's) agammaglobulinemia is not associated with a notable predisposition toward any gastrointestinal illness.⁵⁸ A workup for immunodeficiency is not indicated in patients with giardiasis.⁶⁰

HLA phenotypes—A1/A2 and B12/B27—occur with a higher than expected frequency in patients with giardiasis.⁶¹ T-cell deficiencies in animals are not necessarily risk factors; certain strains of mice are more susceptible to *Giardia muris* infections, however, as are congenitally athymic animals.⁵⁹ In humans the relationship is less clear. With AIDS, for example, the relationship is complex. A history of giardiasis is associated among homosexual men with an odds ratio of over 10 with the development of seropositivity to the human immunodeficiency virus and AIDS.⁶² The incidence of giardiasis among uninfected AIDS patients appears not to be increased, however; homosexual AIDS patients with giardiasis have lower antibody levels than do heterosexual patients with giardiasis—but higher than healthy persons.⁶³

Malnutrition is a likely risk factor for giardiasis, although the cause-and-effect relationship between the two is hard to clarify. In Bangladesh 57% of a malnourished cohort aged 1 to 5 was infected with *G lamblia*.⁶⁴ A Guatemalan study found, among infected infants, reductions in weight gain (but not height) during the second year of life.²³ A Zimbabwean study also correlated infection with *Giardia* species with indices of malnutrition including wasting and stunted growth.⁶⁵ The extent to which immunodeficiency is a significant predisposing cofactor in malnourished patients is unknown.

Persons with blood group A appear to be prone to the recurrence of giardiasis.⁶⁶ This is thought to be related to the antigenic similarities between blood group antigens and *Giardia* species: the organism does not evoke the same immunologic reaction it would if the blood group's glycoproteins differed. Data from Finland showed no correlation between blood groups and (nonrecurrent) giardiasis.⁶⁷ An association between giardiasis and achlorhydria has also been postulated,⁶⁷ although it is essential to control for blood groups because the A blood group is more common among achlorhydric patients.⁶⁸ Cystic fibrosis may be a risk factor: a case-control study (using household family members of cystic fibrosis cases as controls) reported higher infection rates among the patients with cystic fibrosis.⁶⁹

In summary, major host risk factors associated with giardiasis are age and perhaps immunodeficiency. Concomitant infectious diseases may reflect acquisition from a common source. Immunotype, blood type, and achlorhydria share a minor role in contributing a propensity toward infection.

Clinical Symptoms

Most giardial infections, particularly those of infancy, are well tolerated and asymptomatic. When overt illness occurs, the spectrum is wide and ranges from an acute, mild, self-limited gastroenteritis to a protracted (the average duration of diarrhea in one Aspen outbreak was 44 days⁷) and debilitated malabsorptive state due to chronic gastroenteritis.⁷⁰⁻⁷⁵ Frequent symptoms that accompany the gastroenteritis are fatigue, weight loss, abdominal cramps, nausea, and greasy stools.⁶ In children malaise,

dyspeptic disorders, and a failure to thrive may be due to giardiasis, and in adults symptoms may include a diminished capacity for work.²⁶ In the developing world, more than 40% of symptomatic infections last between four and six weeks.²³ The relative proportions of symptomatic and asymptomatic infections in areas with high endemicity are unknown (one study in an outbreak setting¹⁶ documented a high rate of asymptomatic infection).

The small intestine, while the most common, is not the only site of disease. Malabsorptive complications associated with giardial infections include vitamin B₁₂ deficiency due to nutrient competition,⁷⁶ iron deficiency due to impaired absorption,⁷⁷ and hypovitaminosis A.⁷⁸ Less common, documented complications (typically, case reports) include cholecystitis and mechanical jaundice as well as pancreatitis, upper gastrointestinal bleeding, and mesenteric adenitis.⁷⁹⁻⁸¹ Erythema nodosum has been associated with giardiasis,⁸² as have ocular inflammation (iridocyclitis, retinal disease),⁸³ arthritis,⁸⁴ and even psychiatric manifestations in childhood.⁸⁵ Chronic urticaria is reported to occur with giardiasis and subsides with treatment of the infection; the basis is thought to be mucosal injury with the absorption of toxic materials.⁸⁶ Immunoglobulin E levels do not increase in giardiasis.⁸⁷

Pathogenesis

There is no one accepted theory about how *G lamblia* causes disease. Pathologic findings vary but include blunting of small intestinal villi, lymphocytic infiltration, and occasional ulcerations⁸⁸⁻⁹⁰ and do not correlate well with the disease state.⁹¹ Postulated theories regarding pathogenesis include mechanical obstruction by giardial cysts; mucosal damage by nonmechanical means, toxic or metabolic (a metabolic example is the induction of brush border enzyme deficiencies); and deconjugation of bile salts, either by bacterial overgrowth or by giardial organisms.^{59,92} Mucosal invasion is well documented histologically.⁹³ Indian studies have shown that the size of the inoculum of *Giardia* organisms has no effect, over a twofold log range, on either glucose or amino acid transport in intestinal brush border membrane vesicles of infected mice; hence, mechanical obstruction may not be a primary factor in pathogenesis.⁹⁴ In another study, brush border enzyme deficiencies, especially disaccharidase deficiencies, were prominent in a dexamethasone-immunosuppressed mice model.⁹⁵ Not all humans show mucosal damage, and enzyme deficiencies may be a consequence rather than a cause of mucosal disturbances. Bacterial overgrowth is also an inconstant finding.⁹²

The importance of surface lectins has recently been delineated: *G lamblia* produces a lectin activated by human duodenal secretions (a duodenal protease) and inhibited by trypsin inhibitors. The lectin is inhibited competitively by mannose-6-phosphate and agglutinates to the same site in the duodenum where *G lamblia* attaches. The mechanism of lectin-induced organism attachment and resultant disease is unknown. This system is considered analogous to myxovirus activation by proteolytic cleavage of surface viral glycoproteins or the enhancement of *Trypanosoma cruzi* infections by the proteolysis of its surface membrane proteins.⁹⁶

Serum antibodies against a major 82,000-dalton antigen of *Giardia* trophozoites are present among persons with persistent giardiasis.⁹⁷ Substantial indirect fluorescent antibody (IFA) titers have been detected in more than 85% of a Scandinavian adult population, with a fall in titers with treatment.⁹⁸ Indirect fluorescent antibody titers were present as long as 18 months in chronically infected persons in a study from Cincinnati; Indochinese refugees and male

homosexuals with AIDS had higher antibody levels than the uninfected.⁶³ Much of the detected antibody response in populations with high seroprevalence may be due to cross-reaction with other immunogens.⁹⁸ High concentrations of anti-*Giardia* antibodies have also been found in the breast milk of mothers in Bangladesh⁶⁴; these antibodies may act in conjunction with free fatty acids of milk because fatty acids are known to be lethal to both *Giardia* cysts and other parasites.^{99,100}

Ultrastructural studies have elucidated *Giardia* species motility and attachment.¹⁰¹⁻¹⁰³ The trophozoite attaches to the mucosa with a ventral disc whose ultrastructure consists of multiple microtubules, the outer surface of which contains bridges from which emanate microribbons. Motility is achieved by flagellae whose placement determines the direction of movement: posterior flagellae clear a concave space on the undersurface of the organism for purposes of attachment while caudal flagellae alter the movement of substances within the cytoplasm. Flagellae emerge from axonemes, organelles consisting, like the ventral disc, of microtubules but arranged in a geometrically constant pattern: nine peripheral pairs about two central pairs. Organelles degrade on encystation of the organism. The discovery that chitin is a structural component of cysts¹⁰⁴ suggests a possible therapeutic intervention with chitinases.

Diagnosis

A diagnostic workup includes the collection of several stool specimens—typically three—in polyvinyl alcohol fixative (vials should be given to patients with instructions to take them to a laboratory as soon as possible) followed by the identification of trophozoites or cysts by experienced laboratory personnel.¹⁰⁵ A small bowel biopsy or duodenal aspiration facilitates establishing the diagnosis. Combining stool cultures and duodenal aspiration is especially effective in diagnosis,¹⁰⁶ although the string test (Enterotest; Health Development Corporation) may be preferable. A string embedded in a gel capsule is swallowed, with the proximal end of the string taped to the face; the string is removed three to four hours later, when ostensibly it has reached the terminal duodenum; the distal section should contain bile-stained mucus in which motile trophozoites can be detected.^{107,108} An intravenous secretin challenge increases the yield of organisms.¹⁰⁹ A stool enzyme-linked immunosorbent assay is currently available, is 92% to 98% sensitive, nearly 100% specific, and inexpensive; it may eventually replace multiple stool specimens and duodenal aspiration¹¹⁰ and enable a diagnosis using a minimum of technology in a short interval (less than three hours).¹¹¹ Serum IFA levels are commonly present in patients with chronic infection,^{63,98} although the correlation between antibody levels and stool specimen positivity in one malnourished cohort was poor.¹¹² A serum ELISA using cyst antigens has been developed¹¹³ that has high sensitivity (91/92 asymptomatic carriers detected), but specificity is poor (cross-reactivity to heterologous parasitic antigens occurred).

Treatment

Approaches to therapy should consider both the individual and the community: Are individual cases associated with an outbreak? Are physicians receiving current recommendations during an outbreak? It should first be recognized that many infections clear in the absence of treatment.^{73,114} The two most popular agents are nitroimidazoles and quinacrine. The nitroimidazoles include metronidazole, tinidazole, ornidazole, and nimorazole (only the first is marketed in the United States). They are effectively given as

a single dose (500 mg),²⁶ although it is generally recommended that metronidazole be given over a seven- to ten-day course at 500 to 750 mg per day in two to three doses.^{107,115-122} A major concern with metronidazole use is mutagenicity and carcinogenicity. Its use is usually well tolerated, although patients should be reminded of disulfiram-like effects—which may occur as late as 48 hours after therapy—when taken in conjunction with alcohol. Less commonly reported symptoms include headaches, an unpleasant taste in the mouth, stomatitis, paresthesias, dizziness, and transient neutropenia.¹¹⁶ Quinacrine is cheaper, but side effects that may limit compliance¹¹⁹ include gastrointestinal disturbances, blood dyscrasias, urticaria, exfoliative dermatitis, yellowing of the skin, and ocular toxicity; toxic psychoses may occur in 1% to 2% of patients,¹²³ and hepatic necrosis is rare. Furazolidone is another alternative. A liquid form is well tolerated by children,¹²⁴ but gastrointestinal side effects occur, hemolysis may occur in those with a glucose-6-phosphate dehydrogenase deficiency, and it is carcinogenic in animals¹¹⁹; the optimal course appears to be ten days.¹²⁵ The relative efficacy of the compounds is debatable (none of the agents cure more than 85% of cases), and both metronidazole^{26,118,120} and quinacrine¹²⁶ have their strong advocates. In an in vitro system using thymidine incorporation, the relative killing efficacy of nitroimidazoles is greater than that of furazolidone, which is greater than that of quinacrine.¹²⁷ The World Health Organization currently recommends using nitroimidazoles.²⁶

The management of severe cases during pregnancy is problematic; no agent is recommended during early pregnancy. The safety of metronidazole use in children and pregnant women is debatable^{107,121}; while some investigators consider it safe to use during pregnancy,¹¹⁶ in the first trimester its use should be restricted.^{117,118} In a recent study of intestinal parasite carriage, including *Giardia* species, among 147 pregnant Thai refugees, there were no complications associated with the parasite infections. The authors consequently advocated withholding all antiparasitic chemotherapy during pregnancy.¹²⁸ On delivery, a concern with immediate treatment using metronidazole is its passage into breast milk.^{107,129}

Community therapy is directed toward prevention.¹³⁻¹⁷ Water sources of *Giardia* species are typically surface water; contaminated water requires either hyperchlorination or the combined methods of coagulation, sedimentation, and flocculation.¹³ Filtration alone is insufficient, and chlorination alone requires levels of 2 to 5 mg per liter, levels that make water unpalatable. Boiling cyst-infected water immediately destroys the capacity to excyst.¹¹ Ozone has potential promise as an inactivating agent.¹³⁰ For hikers and campers who use limited quantities of water, iodination is sufficient,²⁹ although tetraglycine hydroperiodide is safer and easier to use than crystalline iodine.¹³¹

The proper treatment of institutional outbreaks is controversial.^{40,132} It is often recommended that no treatment be given because such infections are usually well tolerated and control measures are hard to execute. Institutional asymptomatic infections, however, may be the source of more insidious or severe infections in the community. Immunoprophylaxis is not available, and chemoprophylaxis will become feasible only with the availability of cheaper, safer medications.

New Areas for Investigations

Topics currently being investigated that should assist in understanding *Giardia* species and its epidemiology include

protozoal enzymes and analyses of zymodenes (the patterns of isoenzyme variation),¹³³ morphometric observations,¹³⁴ animal reservoirs,²⁶ and antigenic characteristics.^{135,136}

Antigenic characteristics assist in the development of serologic assays and immunotherapy.¹³⁵ Antigenic variation among organisms is known to occur both in vitro and in vivo, although the biologic significance of such variation is unknown.^{135,136} In a recent review of vaccine developments, *G lamblia* was listed among the pathogens for which vaccines are unlikely to be developed in the next decade.¹³⁷ Although much work has been produced in the developing world,* there exists a need for epidemiologic studies outlining different modes of spread in the developed versus the developing world.¹³⁹ Other epidemiologic and therapeutic studies are needed to refine the role and assess the feasibility of eradicating infection in day-care centers among asymptomatic cyst passers—who are well recognized to maintain the disease.⁴⁰⁻⁴⁵

Additional studies on pathogenesis will delineate how lectin activation induces disease,⁹⁶ the primary versus the secondary role of brush border enzyme deficiencies,⁹⁵ structural-physiologic correlates of the organelles,¹⁰¹⁻¹⁰³ and the role of chitin in the disease.¹⁰⁴ Improved, available diagnostic assays are needed, in particular immuno-diagnostic techniques that can establish the diagnosis using either serum or stool specimens.^{110,111} Newer forms of chemotherapy are needed as well as assessments of nitroimidazole derivatives,^{116,120,121} confirming reported cure rates of more than 90%.¹²² The role of the immune system needs to be clarified as well. Further studies delineating the role of the immune system are also needed.

In summary, although *Giardia* species has been a neglected organism with poorly understood pathogenic potential, it is now considered an important cause of gastroenteritis as a consequence of contaminated water and food or venereal exposures. In day-care centers, person-to-person spread is common and asymptomatic cyst passers maintain disease. Periodic major waterborne outbreaks are controlled when attention is directed toward water delivery systems. Current areas for study include parasitic enzymes and antigenic characteristics, relationships to states of impaired immunity, new agents to manage community-acquired (including day-care centers) cases, and ranges of animal reservoirs. The development of safe, effective chemoprophylaxis and immunoprophylaxis should improve the control of giardiasis.

*References 21-23, 52-54, 64, 65, 78, 94, 95, 109, 120, 122, 128, 133, 138.

1. Leeuwenhoek A: Letter 34, 4 Nov 1681, to R. Hooke, MS Roy Soc, In Dobell C (Ed): Antony Leeuwenhoek and His Little Animals, 1st Ed. New York, Dover Publications, 1932

2. Bensen W: Bau und Arten der Gattung *lamblia*. Z Hyg Infekt (Leipzig) 1908; 61:109-114

3. Ackers JP: Giardiasis—Basic parasitology. Proc R Soc Trop Med Hyg 1980; 74:427-429

4. Noc F: Observations sur le cycle évolutif de *Lamblia intestinalis*. Bull Soc Pathol Exot (Paris) 1909; 2:93-97

5. Kofoid CA, Swezy O: Mitosis and fission in the active and encysted phases of *Giardia enterica* (Grassi) of man, with a discussion of the method of origin of bilateral symmetry in the polymastigote flagellates. Univ Calif Pub Zool Berkeley 1922; 20:199-227

6. Brodsky RE, Spencer HC, Schultz MG: Giardiasis in American travelers to the Soviet Union. J Infect Dis 1974; 130:319-323

7. Moore GT, Cross WM, McGuire D, et al: Epidemic giardiasis at a ski resort. N Engl J Med 1969; 281:402-407

8. Bingham AK, Meyer EA: *Giardia* excystation can be induced in vitro in acidic solutions. Nature 1979; 277:301-302

9. Gillin FD, Reiner DS, Gault MJ, et al: Encystation and expression of cyst antigens by *Giardia lamblia* in vitro. Science 1987; 235:1040-1043

10. Christie AB: Gastroenteritis I, chap 8, In Infectious Diseases: Epidemiology and Clinical Practice, 4th ed. Edinburgh, Churchill Livingstone, 1987

11. Holtan NR: Giardiasis—A crimp in the life-style of campers, travelers, and others. Postgrad Med 1988; 83:54-56, 59-61

12. Rendtorff RC, Holt CJ: Experimental transmission of human intestinal protozoan parasite—IV. Attempt to transmit *Entamoeba coli* and *Giardia lamblia* cysts by water. Am J Hyg 1954; 60:320-338

13. Dykes AC, Juranek DD, Lorenz RA, et al: Municipal waterborne giardiasis: An epidemiologic investigation—Beavers implicated as a possible reservoir. Ann Intern Med 1980; 92:165-170

14. Weniger BG, Blaser MJ, Gedroze J, et al: An outbreak of waterborne giardiasis associated with heavy water runoff due to warm weather and volcanic ashfall. Am J Public Health 1983; 73:868-872

15. Istre GR, Dunlop TS, Gaspard GB, et al: Waterborne giardiasis at a mountain resort: Evidence for acquired immunity. Am J Public Health 1984; 74:602-604

16. López CE, Dykes AC, Juranek DD, et al: Waterborne giardiasis—A community-wide outbreak of disease and a high rate of asymptomatic infection. Am J Epidemiol 1981; 112:495-507

17. Jephcott AE, Begg NT, Baker IA: Outbreak of giardiasis associated with mains water in the United Kingdom. Lancet 1986; 1:730-732

18. López CE, Juranek DD, Sinclair SP, et al: Giardiasis in American travelers to Madeira Island, Portugal. Am J Trop Med Hyg 1978; 27:1128-1132

19. Allison DJ: Giardiasis—A recent investigation. Can J Public Health 1984; 75:318-320

20. Walker JC, Conner G, Christopher PJ, et al: A presumed epidemic of giardiasis. Med J Austr 1986; 145:548-549

21. Gilman RH, Brown KH, Visvesvara GS, et al: Epidemiology and serology of *Giardia lamblia* in a developing country—Bangladesh. Trans R Soc Trop Med Hyg 1985; 79:469-473

22. Oyerinde JPO, Ogunbi O, Alonge AA: Age and sex distribution of infections with *Entamoeba histolytica* and *Giardia intestinalis* in the Lagos population. Int J Epidemiol 1977; 6:231-234

23. Farthing MJG, Mata L, Urrutia JJ, et al: Natural history of *Giardia* infection of infants and children in rural Guatemala and its impact on physical growth. Am J Clin Nutr 1986; 43:395-405

24. Gilman RH, Marquis GS, Miranda E, et al: Rapid reinfection by *Giardia lamblia* after treatment in a hyperendemic Third World community. Lancet 1988; 1:343-345

25. Prevention and control of intestinal parasitic infections. WHO Tech Rep Ser 749, 1987

26. Intestinal protozoan and helminthic infections. WHO Tech Rep Ser 666, 1981

27. Hopkins RS, Shillam P, Gaspard B, et al: Waterborne disease in Colorado—Three years' surveillance and 18 outbreaks. Am J Public Health 1985; 75:254-257

28. St Louis ME: Water-related disease outbreaks, 1985. MMWR CDC Surveil Summ 1988; 37:15-24

29. Barbour AG, Nichols CR, Fukushima T: An outbreak of giardiasis in a group of campers. Am J Trop Med Hyg 1976; 25:385-389

30. Porter JD, Ragazzoni HP, Buchanan JD, et al: *Giardia* transmission in a swimming pool. Am J Public Health 1988; 78:659-662

31. Greensmith CT, Stanwick RS, Elliot BE, et al: Giardiasis associated with the use of a water slide. Pediatr Infect Dis J 1988; 7:91-94

32. Osterholm MT, Porfang JC, Ristinen TL, et al: An outbreak of foodborne giardiasis. N Engl J Med 1981; 304:24-28

33. Petersen LR, Carter ML, Hadler JL: A food-borne outbreak of *Giardia lamblia*. J Infect Dis 1988; 157:846-848

34. Petersen K: Ergebnisse von protozoologischen Stuhluntersuchungen 1980 bis 1985 im DDR-Bezirk Rostock. Angew Parasitol 1988; 29:3-10

35. Chattopadhyay B, Fricker E, Gelia CB: Incidence of parasitic infestations in minority group travelers to and new immigrants arriving from the Third World countries. Public Health 1988; 102:245-250

36. Kean BH, William DC, Luminas SK: Epidemic of amoebiasis and giardiasis in a biased population. Br J Vener Dis 1979; 55:375-378

37. Meyers JD, Kuharic HA, Holmes KK: *Giardia lamblia* infection in homosexual men. Br J Vener Dis 1977; 53:54-55

38. Cooper RF, Allison B, Bowles CC, et al: Giardiasis—A common, sexually transmissible parasitic diarrhea with pitfalls in diagnosis. South Med J 1983; 76:863-865

39. Phillips SC, Mildvan D, William DC, et al: Sexual transmission of enteric protozoa and helminths in a venereal-disease clinic population. N Engl J Med 1981; 305:603-606

40. Pickering LK, Woodward WE, DuPont HL, et al: Occurrence of *Giardia lamblia* in children in day care centers. J Pediatr 1984; 104:522-526

41. Black RE, Dykes AC, Sinclair SP, et al: Giardiasis in day-care centers—Evidence of person-to-person transmission. Pediatrics 1977; 60:486-491

42. Boreham PFL, Shepherd RW: Giardiasis in child-care centers (Letter). Med J Aust 1984; 141:263

43. Pickering LK, Evans DG, DuPont HL, et al: Diarrhea caused by *Shigella*, rotavirus, and *Giardia* in day-care centers: Prospective study. J Pediatr 1981; 99:51-56

44. Bartlett AV, Moore M, Gary GW, et al: Diarrheal illness among infants and toddlers in day care centers—I. Epidemiology and pathogens. J Pediatr 1985; 107:495-502

45. Polis MA, Tuazon CU, Alling DW, et al: Transmission of *Giardia lamblia* from a day care center to the community. Am J Public Health 1986; 76:1142-1144

46. Desowitz RS, Wiebenga NH: A survey for intestinal parasites in Oahu school children. Hawaii Med J 1975; 34:21-23

47. Fontaine JP, Delage A, Lauraire MC: Sur l'épidémiologie de la giardiasis. Ann Parasitol Hum Comp 1984; 59:541-554

48. Woo PTK, Paterson WB: *Giardia lamblia* in children in day-care centers in southern Ontario, Canada, and susceptibility of animals to *G lamblia*. Trans R Soc Trop Med Hyg 1986; 80:56-59

49. Harter L, Frost F, Jakubowski W: *Giardia* prevalence among 1 to 3 year old children in two Washington State counties. Am J Public Health 1982; 72:386-388

50. Weiner D, Brooke MM, Witkon A: Investigation of parasitic infections in the central areas of Philadelphia. Am J Trop Med Hyg 1959; 8:625-629

51. Boreham PFL, Phillips RE: Giardiasis in Mount Isa, northwest Queensland. Med J Aust 1986; 144:524-528

52. Peltola H, Kataja M, Mohamed ON, et al: Intestinal parasitism of children and mothers in rural Somalia. Pediatr Infect Dis J 1988; 7:488-492

53. Moore HA, de la Cruz E, Vargas-Mendez O, et al: Diarrheal disease in Costa Rica—II. The prevalence of certain enteric organisms and their relationships to diarrhea. Am J Public Health 1966; 56:442-451

54. Sullivan PS, DuPont HL, Arafat RR, et al: Illness and reservoirs associ-

ated with *Giardia lamblia* infection in rural Egypt—The case against treatment in developing world environments of high endemicity. *Am J Epidemiol* 1988; 127:1272-1281

55. Ament ME, Ochs HD, Davis SD: Structure and function of the gastrointestinal tract in primary immunodeficiency syndrome—A study of 39 patients. *Medicine (Baltimore)* 1973; 52:227-248
56. Janoff EN, Smith PD, Blaser MJ: Acute antibody responses to *Giardia lamblia* are depressed in patients with AIDS. *J Infect Dis* 1988; 157:798-804
57. Webster AD: Giardiasis and immunodeficiency diseases. *Trans R Soc Trop Med Hyg* 1980; 74:440-443
58. Ament ME, Rubin CE: Relation of giardiasis to abnormal intestinal structure and function in gastrointestinal immunodeficiency syndromes. *Gastroenterology* 1972 Feb; 62:216-226
59. Gillon J: Giardiasis—Review of epidemiology, pathogenetic mechanisms, and host responses. *Q J Med* 1984; 53:29-39
60. Ruskin J: Parasitic diseases in compromised hosts. In Rubin RH, Young LS (Eds): *Clinical Approach to Infection in the Compromised Host*. New York, Plenum Books, 1981
61. Roberts-Thomson IC, Mitchell GF, Anders RF, et al: Genetic studies in human and murine giardiasis. *Gut* 1980; 21:397-401
62. Moss AR, Osmond D, Bacchetti P, et al: Risk factors for AIDS and HIV seropositivity in homosexual men. *Am J Epidemiol* 1987; 125:1035-1047
63. Sullivan R, Linneman CC, Clark CS, et al: Seroepidemiologic study of giardiasis patients and high-risk groups in a midwestern city in the United States. *Am J Public Health* 1987; 77:960-963
64. Islam A, Stoll BJ, Ljungström I, et al: *Giardia lamblia* infections in a cohort of Bangladeshi mothers and infants followed for one year. *J Pediatr* 1983; 103:996-1000
65. Loewenson R, Mason PR, Patterson BA: Giardiasis and the nutritional states of Zimbabwean schoolchildren. *Ann Trop Paediatr* 1986; 6:73-78
66. Sotto A, Cabrera S, Castro J, et al: Blood groups in recurrent giardiasis (Letter). *Lancet* 1983; 2:1312-1313
67. Jokipii L, Jokipii AMM: Is predisposition to giardiasis associated with the ABO blood groups? *Am J Trop Med Hyg* 1980; 29:5-7
68. Paulsen O: Blood-group A and giardiasis (Letter). *Lancet* 1977; 2:984
69. Roberts DM, Craft JC, Mather FJ, et al: Prevalence of giardiasis in patients with cystic fibrosis. *J Pediatr* 1988; 112:555-559
70. Heap T: Giardiasis—A common cause of prolonged diarrhoea in adults. *Med J Aust* 1974; 2:592-595
71. Sealy DP, Schuman SH: Endemic giardiasis and day care. *Pediatrics* 1983; 72:154-158
72. Craft JC: *Giardia* and giardiasis in childhood. *Pediatr Infect Dis* 1982; 1:196-211
73. Meyer EA, Radulescu S: *Giardia* and giardiasis. *Adv Parasitol* 1979; 17:1-47
74. Turner JA: Giardiasis and infections with *Dientamoeba fragilis*. *Pediatr Clin North Am* 1985; 32:865-880
75. Wolfe MS: Giardiasis. *N Engl J Med* 1978; 298:319-321
76. Cordingley FT, Crawford GPM: *Giardia* infection causes vitamin B12 deficiency: First case report. *Aust NZ J Med* 1986; 16:78-79
77. De Vizia B, Poggi V, Vajro P, et al: Iron malabsorption in giardiasis. *J Pediatr* 1985; 107:75-78
78. Mandour AM, Abdel-Rahman AM, Hassan HA, et al: *Giardia lamblia* as a cause of malabsorption and hypovitaminosis 'A' in children. *J Egypt Soc Parasitol* 1984; 14:203-206
79. Goldstein F, Thornton JJ, Szydlowski T: Biliary tract dysfunction in giardiasis. *Am J Dig Dis* 1978; 23:559-560
80. Rey C, Escibano JC, Fox M, et al: Mesenteric adenitis secondary to *Giardia lamblia*. *Dig Dis Sci* 1981; 25:968-971
81. Larrosa-Haro A, Coello-Ramirez P: Duodenitis and bleeding of the upper digestive tract in children with giardiasis. *Bol Med Hosp Infant Mex* 1984; 41:674-676
82. Giordano N, Fioravanti A, Mariani A, et al: Erythema nodosum and *Giardia intestinalis*. *Clin Rheumatol* 1985; 4:481-483
83. Anderson ML, Griffith DG: Intestinal giardiasis associated with ocular inflammation. *J Clin Gastroenterol* 1985; 7:169-172
84. Goobar JP: Joint symptoms in giardiasis (Letter). *Lancet* 1977; 1:1010-1011
85. Khan AA: Psychiatric aspects of giardiasis. *Br Med J* 1974; 4:407
86. Valdés Sánchez AF, Cárdenas Oviedo SO, Gómez Echeverría AH: Urticaria crónica: Etiología y diagnóstico. *Alergia* 1985; 32:41-44
87. Geller M, Geller M, Flaherty DK, et al: Serum IgE levels in giardiasis. *Clin Allergy* 1978; 8:69-71
88. Klima M, Gyorkey P, Min K-W, et al: Electron microscopy in the diagnosis of giardiasis. *Arch Pathol Lab Med* 1977; 101:133-135
89. Yardley JH, Takano J, Hendrix TR: Epithelial and other mucosal lesions of the jejunum in giardiasis. *Bull Johns Hopkins Hosp* 1964; 115:389-406
90. Yardley JH, Bayless TM: Giardiasis. *Gastroenterology* 1967; 52:301-304
91. Judd R, Deckelbaum RJ, Weizman Z, et al: Giardiasis in childhood: Poor clinical and histological correlations. *Israel J Med Sci* 1983; 19:818-823
92. Tandon BN, Tandon RK, Satpathy BK, et al: Mechanism of malabsorption in giardiasis: A study of bacterial flora and bile salt deconjugation in upper jejunum. *Gut* 1977; 18:176-181
93. Brandborg LL, Tankersley CB, Gottlieb S, et al: Histological demonstration of mucosal invasion by *Giardia lamblia* in man. *Gastroenterology* 1967; 52:143-150
94. Samra HK, Garg UC, Ganguly NK, et al: Effect of different *Giardia lamblia* inocula on glucose and amino acids transport in the intestinal brush border membrane vesicles of infected mice. *Ann Trop Med Parasitol* 1987; 81:367-372
95. Khanna R, Vinayak VK, Mehta S, et al: *Giardia lamblia* infection in immunosuppressed animals causes severe alterations to brush border membrane enzymes. *Dig Dis Sci* 1988; 33:1147-1152
96. Lev B, Ward H, Keusch GT, et al: Lectin activation in *Giardia lamblia* by host protease—A novel host-parasite interaction. *Science* 1986; 232:71-73
97. Kumkum, Khanna R, Nain CK, et al: Depressed humoral immune responses to surface antigens of *Giardia lamblia* in persistent giardiasis. *Pediatr Infect Dis J* 1988; 7:492-498
98. Jokipii L, Miettinen A, Jokipii AMM: Antibodies to cysts of *Giardia lamblia* in primary giardiasis and in the absence of giardiasis. *J Clin Microbiol* 1988; 26:121-125

99. Gillin FD, Reiner DS, Wang CS: Killing of *Giardia lamblia* trophozoites by normal human milk. *J Cell Biochem* 1983; 23:47-56
100. Hernell O, Ward H, Blackberg L, et al: Killing of *Giardia lamblia* by human milk lipases: An effect mediated by lipolysis of milk lipids. *J Infect Dis* 1986; 153:715-720
101. Crossley R, Holberton DV: Characterization of proteins from the cytoskeleton of *Giardia lamblia*. *J Cell Sci* 1983; 59:81-103
102. Luchtell DL, Lawrence WP, DeWalle FB: Electron microscopy of *Giardia lamblia* cysts. *Appl Environ Microbiol* 1980; 40:821-832
103. Owen RL: The ultrastructural basis of *Giardia* function. *Trans R Soc Trop Med Hyg* 1980; 74:429-433
104. Ward HD, Alroy J, Lev BI, et al: Identification of chitin as a structural component of *Giardia* cysts. *Infect Immun* 1985; 49:629-634
105. Markell EK, Quinn PM: Comparison of immediate polyvinyl alcohol (PVA) fixation with delayed Schaudinn's fixation for the demonstration of protozoa in stool specimens. *Am J Trop Med Hyg* 1977; 26:1139-1142
106. Kamath KR, Murugasu R: A comparative study of four methods of detecting *Giardia lamblia* in children with diarrhoeal disease and malabsorption. *Gastroenterology* 1974; 66:16-21
107. Stevens DP, Mahmoud AAF: Giardiasis: The rediscovery of an ancient pathogen. In Remington J, Swartz M (Eds): *Current Concepts in Infectious Diseases*, 1st Vol. New York, McGraw Hill, 1980
108. Bezjak B: Evaluation of a new technique for sampling duodenal contents in parasitologic diagnosis. *Am J Dig Dis* 1972; 17:845-850
109. Sagaró E, Blanco E, Frago T, et al: Duodenal intubation with secretin stimulus for diagnosis of giardiasis. *Arch Dis Child* 1977; 52:505-507
110. Ungar BLP, Yolken RH, Nash TE, et al: Enzyme-linked immunosorbent assay for the detection of *Giardia lamblia* in fecal specimens. *J Infect Dis* 1984; 149:90-97
111. Green EL, Miles MA, Warhurst DC: Immunodiagnostic detection of *Giardia* antigen in faeces by a rapid visual enzyme-linked immunosorbent assay. *Lancet* 1985; 2:691-693
112. Aubry P: Syndrome de malabsorption et parasitoses intestinales. *Med Trop (Mars)* 1983; 43:317-324
113. Haralabidis ST: Immunodiagnosis of giardiasis by ELISA and studies on cross-reactivity between the anti-*Giardia lamblia* antibodies and some heterologous parasitic antigens and fractions. *Ann Trop Med Parasitol* 1984; 78:295-300
114. Sagi EF, Shapiro M, Deckelbaum RJ: *Giardia lamblia* prevalence, influence on growth, symptomatology in healthy nursery children. *Israel J Med Sci* 1983; 19:815-817
115. Rossignol JF, Maisonneuve H, Cho YW: Nitroimidazoles in the treatment of trichomoniasis, giardiasis, and amebiasis. *Int J Clin Pharmacol Ther Toxicol* 1984; 22:63-72
116. Webster LT: Drugs used in the chemotherapy of protozoal infections, chap 46. In Gilman AG, Goodman LS, Rall TW, et al (Eds): *Goodman and Gilman's The Pharmacological Basis of Therapeutics*. New York, Macmillan, 1985, pp 1052-1054
117. Rosenblatt JE, Edson RS: Symposium on antimicrobial agents—Metronidazole. *Mayo Clin Proc* 1987; 62:1013-1017
118. Goldman P: Drug therapy—Metronidazole. *N Engl J Med* 1980; 303:1212-1218
119. Davidson RA: Issues in clinical parasitology: The treatment of giardiasis. *Am J Gastroenterol* 1984; 79:256-261
120. Speelman P: Single-dose tinidazole for the treatment of giardiasis. *Antimicrob Agents Chemother* 1985; 27:227-229
121. Lindmark DG, Mueller M: Antitrichomonad action, mutagenicity, and reduction of metronidazole and other nitroimidazoles. *Antimicrob Agents Chemother* 1976; 10:476-482
122. Bassily S, Farid Z, el-Masry NA, et al: Treatment of intestinal *E histolytica* and *G lamblia* with metronidazole, tinidazole and ornidazole: A comparative study. *J Trop Med Hyg* 1987; 90:9-12
123. Evans RL, Khalid S, Kinney JL: Antimalarial psychosis revisited. *Arch Dermatol* 1984; 120:765-767
124. Craft JC, Murphy T, Nelson JD: Furazolidone and quinacrine—Comparative study of therapy of giardiasis in children. *Am J Dis Child* 1981; 135:164-166
125. Murphy TV, Nelson JD: Five versus ten days' therapy with furazolidone for giardiasis. *Am J Dis Child* 1983; 137:267-270
126. Babb RR, Peck OC, Vescin FG: Giardiasis: A cause of travelers' diarrhea. *JAMA* 1971; 217:1359-1361
127. Boreham PF, Phillips RE, Shepherd RW: The sensitivity of *Giardia intestinalis* to drugs in vitro. *J Antimicrob Chemother* 1984; 14:449-461
128. D'Alauro F, Lee RV, Pao IK, et al: Intestinal parasites and pregnancy. *Obstet Gynecol* 1985; 66:639-643
129. Ljungström I, Stoll B, Islam A: *Giardia* infection during pregnancy and lactation (Letter). *Trans R Soc Trop Med Hyg* 1987; 81:161
130. Wickramanayake GB, Rubin AJ, Sprout OJ: Inactivation of *Giardia lamblia* cysts with ozone. *Appl Environ Microbiol* 1984; 48:671-672
131. Zemlyn S, Wilson WW, Hellweg PA: A caution on iodine water purification. *West J Med* 1981; 135:166-167
132. Thacker SB, Kimbal AM, Wolfe M, et al: Parasitic disease—Control in a residential facility for the mentally retarded—Failure of selected isolation procedures. *Am J Public Health* 1981; 71:303-305
133. Baveja UK, Jyoti AS, Kaur M, et al: Isoenzyme studies of *Giardia lamblia* isolated from symptomatic cases. *Aust J Exp Biol Med Sci* 1986; 64:119-126
134. Bertram MA, Meyer EA, Anderson DL, et al: A morphometric comparison of five axenic *Giardia* isolates. *J Parasitol* 1984; 70:530-535
135. Smith PD, Gillin FD, Kaushal NA, et al: Analysis of *Giardia lamblia* from Afghanistan, Puerto Rico, Ecuador, and Oregon. *Infect Immun* 1982; 36:714-719
136. Nash TE: Antigenic variation in *Giardia lamblia*. *Exp Parasitol* 1989; 68:238-241
137. Jordan WS: Impediments to the development of additional vaccines: Vaccines against important diseases that will not be available in the next decade. *Rev Infect Dis* 1989; 11:S603-S612
138. Haralabidis STH: Detection of antibodies against *Candida albicans* in *Giardia lamblia* infected individuals. *Acta Trop (Basel)* 1984; 41:303-304
139. Stevens DP: Selective primary health care: Strategies for control of disease in the developing world—XIX: Giardiasis. *Rev Infect Dis* 1985; 7:530-535